

OptimalHealthSecrets.Com
LIA Lifestyle Instruction Aid

a25. Where do we find phase 2 protein inducers in our diet

By Dr. Victor Jean Ouellette

Modified March 6, 2008

Where do we find Phase 2 protein inducers in our diet?	
These foods help normalise diabetes and help protect us against cancer.	
Kaempferol (a flavonoids)[70]	kale, cauliflower, leek, horseradish [71], gooseberries, strawberries [72]
Flavonoids	blueberries, cranberries, bilberries [73]
Ellagic Acid [74]	strawberries, raspberries/blackberries
Silibinin (a flavolignan)	milk thistle fruit [75]
Curcumin (a polyphenolic)	tumeric [76, 77]
Sulforaphane [78](a metabolite of glucoraphanin)	crucifers, (sprouts of certain broccoli cultivars)
Diallyl Sulfide [79]	garlic
Quersetin [80] (an isoflavone)	onions, bilberries, cranberries, lower levels found in apples, cherries, chokecherries, sea buckthorn berries, currants

References

Burnard H. J. Juurlink, Effects of Dietary Intake of Phase 2 Protein Inducers. Medscape General Medicine 5(4) 2003, <http://www.medscape.com/view/article/464799>

70. Uda Y, Price KR, Williamson G, Rhodes MJ. Induction of the anticarcinogenic marker enzyme, quinone reductase, in murine hepatoma cells in vitro by flavonoids. *Cancer Lett.* 1997;120:213-216. Abstract
71. Herrmann K. Flavonols and flavones in food plants: a review. *J Food Technol.* 1976;11:433-448.
72. Hakkinen SH, Karenlampi SO, Heinonen IM, Mykkanen HM, Torronen AR. Content of the flavonols quercetin, myricetin, and kaempferol in 25 edible berries. *J Agric Food Chem.* 1999;47:2274-2279. Abstract
73. Bomser J, Madhavi DL, Singletary K, Smith MA. In vitro anticancer activity of fruit extracts from *Vaccinium* species. *Planta Med.* 1996;62:212-216. Abstract
74. Barch DH, Rundhaugen LM. Ellagic acid induces NAD(P)H:quinone reductase through activation of the antioxidant responsive element of the rat NAD(P)H:quinone reductase gene. *Carcinogenesis.* 1994;15:2065-2068. Abstract
75. Zhao J, Agarwal R. Tissue distribution of silibinin, the major active constituent of silymarin, in mice and its association with enhancement of phase II enzymes: implications in cancer chemoprevention. *Carcinogenesis.* 1999;20:2101-2108. Abstract
76. Dinkova-Kostova AT, Talalay P. Relation of structure of curcumin analogs to their potencies as inducers of phase 2 detoxification enzymes. *Carcinogenesis.* 1999;20:911-914. Abstract
77. Balogun E, Hoque M, Gong P, et al. Curcumin activates the haem oxygenase-1 gene via regulation of Nrf2 and the antioxidant-responsive element. *Biochem J.* 2003;371 (Pt 3):887-895.
78. Fahey JW, Zhang Y, Talalay P. Broccoli sprouts: an exceptionally rich source of inducers of enzymes that protect against chemical carcinogens. *Proc Natl Acad Sci USA.* 1997;94:10367-10372. Abstract
79. Munday R, Munday CM. Relative activities of organosulfur compounds derived from onions and garlic in increasing tissue activities of quinone reductase and glutathione transferase in rat tissues. *Nutr Cancer.* 2001 ;40:205-210. Abstract
80. Valerio LG Jr, Kepa JK, Pickwell GV, Quattrochi LC. Induction of human NAD(P)H:quinone oxidoreductase (NQ01) gene expression by the flavonol quercetin. *Toxicol Lett.* 2001 ;119:49-57. Abstract